

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Comprehensive and Advanced Pharmacology

Journal homepage: <https://www.ijcap.in/>

Review Article

Pharmacognostic study of zingiber officinale

Ambre SG^{1*}, Gaikar VS¹, Sharmale MN¹, Mhaske PB¹, Phalke PL¹¹Matoshri Radha College of Pharmacy, Ahmednagar, Maharashtra, India

ARTICLE INFO

Article history:

Received 23-04-2023

Accepted 20-06-2023

Available online 02-11-2023

Keywords:

Ginger

antioxidant

anti-inflammatory

antipyretic

ABSTRACT

Ginger is scientifically known as *Zingiber officinale*. *Zingiber officinale* {ginger} widely used in pharmaceutical approach. 6-shogaol, 6-paradol, 6-gingerole, zerbubone and zingerole are some active ingredients available in ginger. Ginger has been used commonly to treat diarrhea, indigestion, nausea, heart burn, cough, bronchitis, asthma, diabetes and menstrual irregularities. It consist of 80% moisture, 2% protein, 2% fiber, 1% mineral, 0.9% fat, 12% carbohydrate, volatile oil 1.9%. Volatile oil in ginger contain a-Pinene, camphene, o-Phellandrene, zingiberene, o-bisabolene, geranial.

Ginger mostly preferred warm, humid climate for cultivation. It can be grown both under rain fed and irrigated conditions. Ayurveda literature preferred administration of ginger in both of communicable and non-communicable diseases.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Ginger is subtropical plant grown for its root, ginger rhizomes can be white, yellow and red in colour depend on its variety.¹ The top producing countries of ginger are India, China, Fiji, Indonesia and Peru. In India and china fresh ginger use as flavouring agent in preparation of vegetables and meat products.² Ginger used for treatment of many diseases such as arthritis and rheumatism, indigestion, constipation, and ulcer, atherosclerosis and hypertension, vomiting, cancer. Digestive system is important parts of the body.³ The digest tract including oral cavity, esophagus, stomach, small and large intestine, rectum and finally end with.⁴ Due to GI track infection, smoking, drinking, gender, age, fatty diet The GI cancer is occurs.⁵ It can cause mild side effects including heartburn, diarrhea, burping, and general stomach discomfort.⁶ In western cuisine, ginger is mostly used in sweet application such as ginger ale, ginger bread, ginger snaps, parkin.⁷




Figure 1:

1.1. Taxonomy, distribution, and botanical description

1. Kingdom: Plantae
2. Division: Magnoliophyta

* Corresponding author.

E-mail address: falkepallavi@gmail.com (Ambre SG).

3. Class: Liliopsida
4. Order: Zingiberales
5. Family: Zingiberaceae
6. Genus: Zingiber
7. Species: Zingiber officinale var. Roscoe



Figure 2: Properties of ginger

1.2. Organoleptic properties

1. Taste: Bitter
2. Shape: Nodule
3. Colour: light brown
4. Odour: Characteristic
5. Size: Rhizomes are 2.5 -7.5 cm in length
6. Fractura: Easy and splintery
7. Surface: Irregular

1.3. Physicochemical properties

1. Ash: 1.02 g/100g
2. Moisture: 98.23 g/100g
3. Protein: 0.02 g/100g
4. Crude fiber: 0.00g/100g
5. Carbohydrate: 0.22g/100g
6. Energy: 5.40g/100g

1.4. Solubility

Soluble: chloroform, carbon tetra chloride, diethyl ether, n-hexane, 100% alcohol
Insoluble: water.⁸

1.5. Therapeutic properties of ginger

1. Antioxidant Activity
2. Antifungal Activity
3. Antibacterial Activity
4. Anti-inflammatory Activity
5. Anti-Cancer Activity

Chemical constituents:⁹

1. 6-paradol
 - (a) Molecular formula=C₁₇H₂₆O₄
 - (b) Molecular weight =278.392g/mol
2. 6-shogaol¹⁰
 - (a) Molecular formula =C₁₇H₂₄O₃
 - (b) Molecular weight=276.4g/mol
3. 6-gingerol
 - (a) Molecular formula =C₁₇H₂₄O₄
 - (b) Molecular weight =294.4 g/mol
4. Isolation of 6 gingerol
Maximum yield of 6-gingerol was obtained with 80% ethanol as extracting solvent at 30°C. We obtained increased yield (7%) of extraction by pretreatment with ultrasonication.¹¹
5. Zerumbone
 - (a) Molecular formula =C₁₅H₂₂O
 - (b) Molecular weight = 218.3 g/mol

2. Isolation of Zerumbone

Was isolated from fresh rhizomes of Zingiber zerumbet Smith in yields of 0.3-0.4% by simple steam distillation and recrystallization.¹²

2.1. Antioxidant activity

Z. officinale is effective in parkinsons disease. Antioxidant stabilize free radicals before they attack target in biological cell. For energy production generated during the process of oxidation, some free radicals are essential. Increases in the production of free radicals show oxidative stress that can lead to damage to DNA.¹³ It has powerful antioxidant activity due to its oil which has a protective effect on DNA damage.¹⁴ Ginger show antioxidant effect in human chondrocyte cell. The ginger extract showed antioxidant effect in inhibiting DPPH radical, IC₅₀ was 4.25 mg/ml. DPPH analysis is one of the test use to prove the ability of the components of the ginger extract to act as donor of hydrogen atoms.¹⁵ It inhibits an enzyme, namely, xanthine oxidase, which is mainly involved in the generation of reactive oxygen species. Antioxidant activity of the plants is due to the presence of flavones, isoflavones, flavonoids, anthocyanin, coumarin lignans, catechins and isocatechins.¹⁶

2.2. Antifungal Activity

The antifungal principle of ginger are Gingerols and gingerdiol. Ginger powder is effective in several antifungal disease (Ernst and Pittler, 2004; Ramkisson et al 2012).

2.3. Antidiabetic activity

In diabetes, many studies show that ginger and other plants have effective both preventively and therapeutically. Diabetes is major global health problem in worldwide. The cause of diabetes is low insulin.¹⁷ A study show that ginger increase uptake of glucose into muscles cell without using insulin.¹⁸ Ginger also protect a diabetics liver, kidney and central nervous system. In Australia, the university Sydney found ginger was effective in glycemic control for people with type 2 diabetes. In high fat diets the ethanolic extract of ginger was found to reduced total cholesterol, body weight, triglycerides, LDL cholesterol, insulin, glucose, free fatty acids and phospholipid.¹⁹

2.4. Antibacterial activity

Ginger has strong antibacterial properties. Ginger show antibacterial effect on both gram (+ve) and gram(-ve) bacteria. Ginger gives antibacterial activity against *Escherchia coli*, *Salmonella enteriditis* and *Staphylococcus Aureus*. Main components,including –gingerol and²⁰ –gingerol, isolated from ginger rhizome displayed antibacterial activity against periodontal bacteria.

2.5. Anti-Inflammatory activity

Z. officinale is highly effective in inflammations associated with alimentary canal. Ginger is useful in treating inflammation, pain and rheumatism.²¹ Thomson et al.2002, Wigler et al.2003. Ramadan et. Al 2011, Funk et al. 2016. Clinically investigated the safety and effectiveness of ginger.²² Ginger and its derivatives are used in many countries to boost the immune system.{22} In ancient herbs use to support the body's immune response, ginger has the capacity to reduce inflammation, swelling, and discomfort.²³ First group to investigate the effect of ginger on platelet, aggregation by using four ginger extract, produced using different solvent i.e.aqueous, n-hexane, chloroform and ethyl acetate.²⁴

2.6. Anti-cancer

Z. officinale exhibits anti-inflammatory and anti-tumorigenic effects. Ginger is effective in preventing growth of cancer has been studied in a variety of cancer types, including colorectal cancer, breast cancer, lymphoma, hepatoma, skin cancer, liver cancer.²³ Preclinical studies have shown that ginger extract and its constituents posses chemopreventive and antineoplastic properties in gastric cancer.²⁵ A component of Asian ginger, zerumbone, also inhibits growth and proliferation of pancreatic cancer through different mechanisms. In animal model, ginger suppresses ethionine –induced liver carcinogenesis by scavenging the free radical formation and by reducing lipid peroxidation, thus ginger prevents rat

hepatocarcinogenesis.²⁶

2.7. Side Effect

In generally *Z. officinale* is considered safe for use .There is a slight risk of toxicity in normal doses, but some people may have indigestion.¹⁷ Use ginger with caution in people with liver disorder or liver disease who have been exposed before.

2.8. Land preparation

Prepare field by ploughing land two-three times. After ploughing do planking operation. For turmeric planting, beds of 15 cm in height, 1cm width and of convenient length are prepared. Keep distance of 50 cm between beds.²⁷

3. Sowing

3.1. Time of sowing

Sowing is done by first weak of May – June.

3.2. Spacing

Deep plant distance of 15-20 cm between the row and 30 cm between two plant.

3.3. Sowing depth

Depth should be near about 3-4 cm.

3.4. Method of sowing

Direct sowing also transplanting method is used for planting.

3.5. Seed rate

For sowing, select fresh and diseased free rhizomes (mother rhizomes as well as fingers) are used. Seed rate of 480-720 kg is sufficient foe sowing one acre land.

3.6. Seed treatment

Before sowing carry out rhizomes treatment with Mancozeb@3 gm/ltr of water. Dip rhizomes for 30 min in solution. It will protect rhizhomes from fungal infestation. After treatment dry rhizomes in shade for 3-4 hours.

3.7. Harvesting

In 8 months crop is ready for harvesting. For fresh spice purpose, ginger is harvested from 6th month and if it is to be used for processing, then it is harvested after 8 months. Right time for harvesting of ginger is when leaves gets yellow and dry out completely. Remove rhizomes by digging and after harvesting clean rhizomes by thoroughly

washing in water for 2 to 3 times. Then dry them in shades for 2-3 days.

3.8. Post harvest

For dry ginger purpose, only the outer skin is to be peeled and then dried in the sun for a week. Outer skin is removed with help of special knife or split bamboos with pointed end. {38} yield of dry ginger is about 16-25% of green ginger.

3.9. Storage

Fresh disease free rhizomes are selected then treated with solution of carbendazim+macrozeb@40gm/10 ltr of water for 30 min. It will prevent rotting of rhizomes in storage. Then dry the rhizomes in pit of convenient size, covered with plank having 2-3 holes for aeration. But storing rhizomes in pit, spread 1 inch thickness sand layers.




Figure 3: Process: ¹⁹

4. Acknowledge

My sincer gratitude goes to Miss Phalke mam and may be loud parents for their continues support.

5. Source of Funding

None.

6. Conflict of Interest

None.

References

1. Frondoza CG, Sohrabi A, Polotsky A, Phan PV, Hungerford DS, Lindmark L, et al. An in vitro screening assay for inhibitors of proinflammatory mediators in herbal extracts using human synovioocyte cultures. *In Vitro Cell Dev Biol Anim.* 2004;40(3-4):95–101.
2. Jiang H, Xie Z, Koo HJ, Mclaughlin SP, Timmermann BN, Gang DR, et al. Metabolic profiling and phylogenetic analysis of medicinal Zingiber species: Tools for authentication of ginger. *Phytochemistry.* 2006;67(15):1673–85.
3. Bhargava S, Dhabhai K, Batra A, Sharma A, Malhotra B. Zingiber Officinale: Chemical And Phytochemical Screening And Evaluation Of Its Antimicrobial Activities. *J Chem Pharm Res.* 2012;4(1):360–4.
4. Ahmed RS, Suke SG, Seth V, Chakraborti A, Tripathi AK, Banerjee BD, et al. Protective effects of dietary ginger (Zingiber officinale Rosc.) on lindane-induced oxidative stress in rats. *Phytother Res.* 2008;22(7):902–6.
5. Hussein MR, Dief EA, Ghait AAE, Adly MA, Abdelraheem MH. Morphological evaluation of the radioprotective effects of melatonin against X-ray induced early and acute testis damage in Albino rats: an animal model. *Int J Exp Pathol.* 2006;87(3):237–50.
6. Available from: http://hukor.kemkes.go.id/uploads/produk_hukum/KMK_No._HK_01_07-MENKES-187-2017_ttg_Formularium_Ramuan_Obat_Tradisional_Indonesia_.pdf.
7. Lutsey PL, Steffen LM, Stevens J. Dietary intake and the development of the metabolic syndrome: the atherosclerosis risk in communities study. *Circulation.* 2008;117(6):754–61.
8. Forouzan S, Bahmani M, Parsaei P, Mohsenzadegan A, Gholami-Ahangaran M, Sadeghi E, et al. Anti-Parasitic Activities of Zingiber officinale Methanolic Extract on *Limnatis nilotica*. *Global Vet.* 2012;9(2):144–8.
9. Ho SC, Chang KS, Lin CC. Anti-neuroinflammatory capacity of fresh ginger is attributed mainly to 10-gingerol. *Food Chem.* 2013;141(3):3183–91.
10. Bhattarai S, Tran VH, Duke CC. The Stability of Gingerol and Shogaol in Aqueous Solutions. *J Pharm Sci.* 2001;90(10):1658–64.
11. Ghasemzadeh A, Jaafar HZE, Rahmat A. Synthesis of Phenolics and Flavonoids in Ginger (Zingiber officinale Roscoe) and Their Effects on Photosynthesis Rate. *Int J Mol Sci.* 2010;11(11):4539–55.
12. Storlien LH, Baur LA, Kriketos AD. Dietary fats and insulin action. *Diabetologia.* 1996;39(6):621–31.
13. Hoque MA, Bala BK, Hossain MA, Uddin MB. Drying Kinetics Of Ginger Rhizome (Zingiber Officinale). *Bangladesh J Agril Res.* 2013;38(2):301–19.
14. Jolad SD, Lantz RC, Solyom AM, Chengj, Bates RB, Timmermann BN, et al. Fresh organically grown ginger (Zingiber officinale): composition and effects on LPS-induced PGE2 production. *Phytochemistry.* 2004;65(13):1937–54.
15. Afzal M, Al-Hadidi D, Menon M, Pesek J, Dhama MI. Ginger: an ethnomedical, chemical and pharmacological review. *Drug Metabol Drug Interact.* 2001;18(3-4):159–90.
16. Pearson TA, Mensah GA, Alexander RW, Anderson JL, Cannon RO, Criqui M, et al. Markers of inflammation and cardiovascular disease: application to clinical and public health practice: A statement for healthcare professionals from the Centers for Disease Control and Prevention and the American Heart Association. *Circulation.* 2003;107(3):499–511.
17. Garace S, Sankari M, Gopi. Antimicrobial Activity of Ethanolic Extract of Zingiber Officinale – An in vitro Study. *J Pharm Sci Res.* 2017;9(9):1417–9.
18. Yadav AK, Reetu, Garg A. Antidiabetic Effects of [10]-Gingerol in Streptozotocin- and High-Fat Diet-Induced Diabetic Rats. *Asian J Pharm Clin Res.* 2019;12(11):77–80.
19. Andriyani R, Budiati TA, Pudjiraharti S. Total Phenolic Content, Antioxidant and Anti-Bacterial Activity of Zingiberis officinale Rhizome. *Procedia Chem.* 2015;16:149–54.

20. Stewart JJ, Wood MJ, Wood CD, Mims ME. Effects of ginger on motion sickness susceptibility and gastric function. *Pharmacology*. 1991;42(2):111–20.
21. Mcgrath KCY, Li XH, Puranik R. Role of 3beta-hydroxysteroid-delta 24 reductase in mediating antiinflammatory effects of high-density lipoproteins in endothelial cells. *Arterioscler Thromb Vasc Biol*. 2009;29(6):877–82.
22. Ghosh S, May MJ, Kopp EB. NF-kappa B and Rel proteins: evolutionarily conserved mediators of immune responses. *Annu Rev Immunol*. 1998;16:225–60.
23. Oyagbemi AA, Saba AB, Azeez OI. Molecular targetsof [6]-gingerol: its potential roles in cancer chemoprevention. *BioFactors*. 2010;36(3):169–78.
24. Islam MS, Mia M, Apu MI, Halder J, Rahman MF, Islam M, et al. A Comprehensive Review On Region Based Traditional Ayurvedic Practitioner’s Plants Secondary Metabolites And Their Phytochemical Activities In Bangladesh. *J Pharmacogn* . 2015;3(6):202–16.
25. Tripathi S, Maier KG, Bruch D, Kittur DS. Effectof 6-gingerol on pro-inflammatory cytokine production andcostimulatory molecule expression in murine peritoneal macrophages. *J Surg Res*. 2007;138(2):209–13.
26. Lee C, Park GH, Kim C, Jang J. [6]-Gingerol attenuates β -amyloid-induced oxidative cell death via fortifying cellular antioxidant defense system. *Food Chem Toxicol*. 2011;49(6):1261–9.
27. Basciano H, Federico L, Adeli K. Fructose, insulin resistance, and metabolic dyslipidemia. *Nutr Metab*. 2005;2(5). doi:10.1186/1743-7075-2-5.

Author biography

Ambre SG, Student

Gaikar VS, Student

Sharmale MN, Assistant Professor

Mhaske PB, Assistant Professor

Phalke PL, Principal

Cite this article: Ambre SG, Gaikar VS, Sharmale MN, Mhaske PB, Phalke PL. Pharmacognostic study of zingiber officinale. *IP Int J Comprehensive Adv Pharmacol* 2023;8(3):138-142.