

## Pharmacoepidemiological study of the usage of caripill for the dengue patients in an urban health setup among the registered medical practitioners

T Janagan<sup>1</sup>, S.A Sridevi<sup>2\*</sup>

<sup>1,2</sup>Professor, Dept. of Pharmacology, <sup>1</sup>Sree Muthukumaran Medical College and Hospital, Chikkarayapuram, Chennai, <sup>2</sup>ACS Medical College and Hospital, Chennai, Tamil Nadu, India

**\*Corresponding Author: S.A Sridevi**

Email: sridevidakshya@gmail.com

### Abstract

Dengue is a prevailing infectious disease in India currently associated with high mortality and morbidity. It is a viral disease without any specific treatment or specific drug availability. Some reports show that carica papaya leaves extract are said to improve the platelet count among dengue patients: but to support this, the prescription pattern and its common usage among the registered medical practitioners are not available.

**Objectives:** This study is to evaluate the pharmaco epidemiology of the use of caripill among the registered medical practitioners in an urban population, Chennai. By knowing this, the popularity of the drug and its usage in the market can be evaluated.

**Keywords:** Caripill, Dengue, Pharmacoepidemiology.

### Introduction

Dengue is an acute infectious disease caused by mosquito and associated with significant mortality and morbidity.<sup>1</sup> Today about 50% of the world's population, live in areas which are endemic for dengue transmission, and around 125 countries are endemic for the same.<sup>2</sup> Globally, around 50–200 million dengue infection, 500,000 episodes of severe dengue, and 20,000 dengue-related deaths are reported annually.<sup>3</sup> The incidence and prevalence of dengue infection are expected to increase in future because of changes in climate, travel, socioeconomic status, trade, and viral characteristics.<sup>3</sup> There is 40% risk of contracting dengue if one resides in the endemic area and out of these, only 0.5% has risk of developing a serious form of the disease. Clinically, dengue causes moderately high-grade fever, rash with severe aches, mostly severe backache, and pain behind the eyes. Following this, there is an apparent phase of improvement and in severe forms symptoms such as vomiting, pain in the abdomen, red spots, reduction in urine output, change in alertness, and bleeding can occur. The red flag sign, citing the severity in serious dengue is a fluid leak from blood vessels and capillaries from the nose, gums, skin, lung vessels, intra-abdominal areas, etc.<sup>4</sup>

Bleeding of variable severity is seen mainly in the severe form of dengue, i.e., dengue hemorrhagic fever. Pathogenesis of bleeding involves leakage of plasma from the blood vessels. It seems to be due to the destruction of platelets through complement system or depression of bone marrow due to dengue viral infection or both. It is also observed that platelets that escape from the destruction become nonfunctional or less functional.<sup>5,6,7</sup> Direct correlation between platelet count and prognosis of dengue is reported in many published studies.<sup>5,6,8-10</sup>

With the recent outbreak of dengue in India, there has reportedly been a sudden increase in the demand for papaya leaves juice across India. The print media, social network, and national media have started coming up with opinion pieces, news items, and blogs on the use of papaya in the treatment of dengue which was instantly taken up by the community due to its perceived beneficial effect on the platelet count.<sup>11</sup> Literature cites the therapeutic effects of aqueous extract of papaya (*Carica papaya*) leaves that are presumed to be due to several active components (flavonoids, alkaloids, enzymes, and minerals) which may have antioxidant and immunomodulatory effects.<sup>12-15</sup>

**Table 1:**

Clinic with a digital number	Number of dengue fever patients (as per inclusion criteria)	Prescription pattern	
		Conventional pain killers only	Painkillers + caripill tablet
Clinic 1	35	11	24
Clinic 2	12	7	5
Clinic 3	43	16	27

Clinic 4	28	8	20
Clinic 5	54	41	13
Clinic 6	18	11	7
Clinic 7	42	27	15
Clinic 8	39	12	27
Clinic 9	56	26	30
Clinic 10	36	17	19
Clinic 11	24	8	16
Clinic 12	32	13	19

With this above literature on the efficacy of carica papaya extract, the prescription pattern and use of the extract, caripill is assessed among the registered medical practitioners.

### Materials and Methods

It is an observational study based on the data collected from the registered medical practitioners who are doing their general practice in Chennai between June 2019 to November 2019. For this study, the data of the dengue patient found positive and the treatment modalities were collected. Most of the doctors are full time general practitioners with an outpatient clinic setup. The records of the patients included their age, sex, clinical signs and symptoms, the investigations done for evaluating dengue infection (NS1 positive), platelet count and complete hemogram. Those patients whose platelets are normal will be categorized under dengue fever and those patients whose platelets fall below 50000/ $\mu$ l of blood will come under dengue hemorrhagic fever. This was collected for a 6 month period between June 2019 to November 2019.

Here the inclusion criteria will be patients in the age group of 25 to 50 years, both male and female, NS1 positive, dengue fever (i.e. platelet count normal). Exclusion criteria will be patients less than 25 years, more than 50 years, patients with other co

morbid conditions like renal or hepatic failure, pregnancy, hematological disorders like hemophilia, idiopathic thrombocytopenia. So, the prescription pattern for those patients identified with dengue fever alone evaluated.

**Data collection and analysis:** The data was collected from 12 outpatient clinics in an urban area of Chennai, Mogappair. All the doctors running the clinic were registered medical practitioners. The data included in the order given to the clinic in a digital format, number of patients with NS1 positivity and normal blood platelet counts and the prescription pattern which included pain killers like only acetaminophen in some patients and those

with add on therapy of caripill with the dosage were recorded. The data are as follows:

The dosage of caripill is 1100mg thrice a day for five days. This is the drug manufactured by microlabs india.

**Statistical analysis:**

The data given above in the tabular column are analysed statistically. The figure 1 shows the pattern of each clinic showing their prescription pattern with and without caripill. Fig. 1:




Fig. 1

### Results

The efficacy of carica papaya leaf extract is evident in this study.<sup>17</sup> In our study based on the proved efficacy of carica papaya leaf extract, the prescription pattern and usage of the drugs were common among many practitioners. It is evident from the above figure 1 that around 8 out of 12 clinics have prescribed caripill tablet than the conventional pain killer regimen. The tablet was given for a period of five days and they recovered after a week duration.

### Discussion

From this study it is evident that the tablet caripill is very much popular for its efficacy and hence many practitioners are prescribing caripill in dengue fever patients to prevent the progression of the disease to the next level of dengue hemorrhagic fever. This avoids the unwanted inpatient admission of the patients. An addition of a drug improves the platelet count and thereby prevents hospitalization.

## Acknowledgements

Special thanks to all the medical practitioners who cooperated for this study. They were helpful in providing data to look for the pharmaco epidemiological pattern of this drug, caripill.

## Source of Funding

None.

## Conflict of Interest

None.

## References

- Guzmán MG, Kourí G. Dengue: An update. *Lancet Infect Dis* 2002;2:33-42. [PubMed] [Google Scholar]
- Gubler DJ. Dengue, urbanization and globalization: The unholy trinity of the 21st century. *Trop Med Health* 2011;39(4):3-11. [PMC free article] [PubMed] [Google Scholar]
- Murray NE, Quam MB, Wilder-Smith A. Epidemiology of dengue: Past, present and future prospects. *Clin Epidemiol* 2013;5:299-309. [PMC free article] [PubMed] [Google Scholar]
- Ho TS, Wang SM, Lin YS, Liu CC. Clinical and laboratory predictive markers for acute dengue infection. *J Biomed Sci* 2013;20:75. [PMC free article] [PubMed] [Google Scholar]
- Makroo RN, Raina V, Kumar P, Kanth RK. Role of platelet transfusion in the management of dengue patients in a tertiary care hospital. *Asian J Transfus Sci* 2007;1:4-7. [PMC free article] [PubMed] [Google Scholar]
- Krishnamurti C, Kalayanaraj S, Cutting MA, Peat RA, Rothwell SW, Reid TJ, et al. Mechanisms of hemorrhage in dengue without circulatory collapse. *Am J Trop Med Hyg* 2001;65:840-7. [PubMed] [Google Scholar]
- de Azeredo EL, Monteiro RQ, de-Oliveira Pinto LM. Thrombocytopenia in dengue: Interrelationship between virus and the imbalance between coagulation and fibrinolysis and inflammatory mediators. *Med Infl* 2015;2015:313842. [PMC free article] [PubMed] [Google Scholar]
- Jayashree K, Manasa GC, Pallavi P, Manjunath GV. Evaluation of platelets as predictive parameters in dengue fever. *Indian J Hematol Blood Transfus* 2011;27:127-30. [PMC free article] [PubMed] [Google Scholar]
- Noisakran S, Chokephaibulkit K, Songprakhon P, Onlamoon N, Hsiao HM, Villinger F, et al. A re-evaluation of the mechanisms leading to dengue hemorrhagic fever. *Ann N Y Acad Sci* 2009;1171(1):E24-35. [PubMed] [Google Scholar]
- Potts JA, Gibbons RV, Rothman AL, Srikiatkachorn A, Thomas SJ, Supradish PO, et al. Prediction of dengue disease severity among pediatric Thai patients using early clinical laboratory indicators. *PLoS Negl Trop Dis* 2010;4:e769. [PMC free article] [PubMed] [Google Scholar]
- Jha DN. Doctors Unsure but Papaya a Hit for Dengue Treatment. *The Times of India*. 2015 Sep 16; [Google Scholar]
- Sarala N, Paknikar S. Papaya extract to treat dengue: A novel therapeutic option? *Ann Med Health Sci Res* 2014;4:320-4. [PMC free article] [PubMed] [Google Scholar]
- Subenthiran S, Choon TC, Cheong KC, Thayan R, Teck MB, Muniandy PK, et al. *Carica papaya* leaves juice significantly accelerates the rate of increase in platelet count among patients with dengue fever and dengue haemorrhagic fever. *Evid Based Complement Alternat Med* 2013;2013:616737. [PMC free article] [PubMed] [Google Scholar]
- Senthilvel P, Lavanya P, Kumar KM, Swetha R, Anitha P, Bag S, et al. Flavonoid from *Carica papaya* inhibits NS2B-NS3 protease and prevents dengue 2 viral assembly. *Bioinformation*. 2013;9:889-95. [PMC free article] [PubMed] [Google Scholar]
- Dharmarathna SL, Wickramasinghe S, Waduge RN, Rajapakse RP, Kularatne SA. Does *Carica papaya* leaf-extract increase the platelet count? An experimental study in a murine model. *Asian Pac J Trop Biomed* 2013;3:720-4. [PMC free article] [PubMed] [Google Scholar]
- Yunita F, Hanani E, Kristianto J. The effect of *Carica papaya* L. leaves extract capsules on platelets count and hematocrit level in dengue fever patient. *Int J Med Aromat Plants* 2012;2:573-8. [Google Scholar]

**How to cite:** Janagan T, Sridevi SA, Pharmacoepidemiological study of the usage of caripill for the dengue patients in an urban health setup among the registered medical practitioners. *IP Int J Comprehensive Adv Pharmacol* 2020;5(1):42-4.